Identificazione del paziente per prodotto per articolazione

Alberto Migliore

I° Congresso Nazionale Società Italiana G.U.I.D.A.

Per la Gestione Unificata e Interdisciplinare del Dolore Muscolo – scheletrico e dell'Algodistrofia

Napoli, 9 -11 Marzo 2017

Hotel Royal Continental

Le opzioni

- ✓ HA
- ✓ Steroide
- ✓ PRP
- ✓ Collagene

COME AGISCE
L'ACIDO
IALURONICO

Meccanismo di azione dell'acido ialuronico

- L'acido ialuronico si lega diversi recettori specifici
- Tali recettori appartengono alla famiglia delle hyaloderine
- Essi sono: CD44, ICAM1, HARE
- RHAMM (recettore per la mobilità mediata dall'acido ialuronico)
- Il più importante di essi è il CD 44

RECCETTORI PER HA: IALADERINE DI MEMBRANA

Different stages of OA

Same expression of CD44R?

Indicazioni

Terapia di 1° linea

- artrosi di grado iniziale intermedio (K/LI-II-III)
- artrosi di grado elevato nel giovane adulto

Terapia di 2° linea

- artrosi di grado elevato inoperabile o in attesa di intervento

La posologia sintomatica

Anca: 1 iniezione ogni sei mesi, aumentabile fino a 4/anno

Ginocchio: 1 ciclo ogni 8-12 mesi

**Personalizzazione del
trattamento**

Ciclicità del trattamento

Attenzione alla corretta diagnosi !

- Entesite (peritrocanterica, zampa d'oca)
- Sofferenza dell'osso sottocondrale
- Borsite (ileopsoas, cisti di baker)
- Condrocalcinosi
- Coxite in corso di spondiloartrite

ANTIAGE TEP

Differences among Branded Hyaluronic Acids in Italy, Part 1: Data from *In Vitro* and Animal Studies and Instructions for Use

A. Migliore¹, E. Bizzi¹, O. De Lucia², A. Delle Sedie³, M. Bentivegna⁴, Asmaa Mahmoud^{5,6} and C. Foti⁷

¹Operative Unit of Rheumatology, S. Pietro Fatebenefratelli Hospital, Rome, Italy. ²Division and Chair of Rheumatology, Gaetano Pini Institute, Milan, Italy. ³Rheumatology Unit, University of Pisa, Pisa, Italy. ⁴Rheumatology Provincial Network Coordinator, ASP 7, Scicli Hospital, Ragusa, Italy. ⁵Physical Medicine, Rheumatology and Rehabilitation Department, Ain Shams University, Cairo, Egypt. ⁶Physical and Rehabilitation Medicine, Tor Vergata University, Rome, Italy. ⁷Physical and Rehabilitation Medicine, Clinical Sciences and Translational Medicine Department, Tor Vergata University, Rome, Italy.

CLINICAL MEDICINE INSIGHTS: ARTHRITIS AND MUSCULOSKELETAL DISORDERS 2016:9

Differences Regarding Branded HA in Italy, Part 2: Data from Clinical Studies on Knee, Hip, Shoulder, Ankle, Temporomandibular Joint, Vertebral Facets, and Carpometacarpal Joint

A. Migliore¹, E. Bizzi¹, O. De Lucia², A. Delle Sedie³, S. Tropea⁴, M. Bentivegna⁵, A. Mahmoud^{6,7} and C. Foti⁸

CLINICAL MEDICINE INSIGHTS: ARTHRITIS AND MUSCULOSKELETAL DISORDERS 2016:9

67 Acidi Jaluronici commercializzati per uso intra-articolare

Arthrum
Arthrum 2,5%
Artrosulfur HA
Artz®/Supartz®
Candrovisc
Coxarthrum®
Durolane®
Durolane sj
Euflexxa®
Fermathron S
Fermathron plus
Go-On®
Go-On® Mini
Go-On matrix

Hyalart®
Hyalgan®
Hyalubrix®
Hyalubrix® 60/Hyalone®
Hymovis®
Inartral
Intrigel® 0.8%
Intrigel® 1.6%
Intrigel® Mini
Jointex®
Jointex® Starter
Jointex® Mini
Jonexa®

Kartilage
Kartilage cross
MonoVisc®
Orthovisc®
Orthovisc® mini
Ostenil®
Ostenil® Plus
Ostenil® mini
Proial
Promovia
RenehaVis®
Rhizarthrum®
Sinovial®
Sinovial® Forte
Sinovial® Mini
Sinovial® One
SportVis®

Structovial®
Synocrom®
Synocrom® mini
Synocrom® Forte
Synocrom® Forte
One
Synolis® V-A
Synvisc®
Synvisc® One
Yaral®
Yaral® Forte
Yaral® Mini
Viscoplus
Viscoplus gel

Indicazioni alla somministrazione

Seguendo il foglietto illustrativo degli acidi ialuronici:

Controindicazioni alla somministrazione

Contents lists available at [ScienceDirect](#)

Seminars in Arthritis and Rheumatism

journal homepage: www.elsevier.com/locate/semarthrit

Consensus statement on viscosupplementation with hyaluronic acid for the management of osteoarthritis

Yves Henrotin, MD^{a,b}, Raghu Raman, MD^c, Pascal Richette, MD^{d,e}, Hervé Bard, MD^f, Jörg Jerosch, MD^g, Thierry Conrozier, MD^{h,*}, Xavier Chevalier, MDⁱ, Alberto Migliore, MD^j

Semin Arthritis Rheum. 2015 Apr 30

ACCORDO UNANIME

Consensus statement on viscosupplementation with hyaluronic acid for the management of osteoarthritis

Yves Henrotin, MD^{a,b}, Raghu Raman, MD^c, Pascal Richette, MD^{d,e}, Hervé Bard, MD^f, Jörg Jerosch, MD^g, Thierry Conrozier, MD^{h,*}, Xavier Chevalier, MDⁱ, Alberto Migliore, MD^j

La Vs é efficace nel trattamento della OA di ginocchio di grado lieve e moderato

La Vs non é una alternativa alla chirurgia nelle forme avanzate di OA dell'anca

La VS é un trattamento ben tollerato dell' OA del ginocchio e di altre articolazioni

La VS non dovrebbe essere usata solo nei casi di fallimento della terapia analgesica e con NSAID.

La VS va proposta come terapia “positiva” e non come una terapia da prescrivere in “mancanza di nient’altro di meglio”

ACCORDO UNANIME

Consensus statement on viscosupplementation with hyaluronic acid for the management of osteoarthritis

Yves Henrotin, MD^{a,b}, Raghu Raman, MD^c, Pascal Richette, MD^{d,e}, Hervé Bard, MD^f, Jörg Jerosch, MD^g, Thierry Conrozier, MD^{h,*}, Xavier Chevalier, MDⁱ, Alberto Migliore, MD^j

- ✓ Il regime di dosaggio dovrebbe essere deciso in base alla EBM
- ✓ Il cross-linking prolunga la permanenza dell'acido ialuronico nella HA
- ✓ Il migliore approccio per le infiltrazioni di ginocchio é il laterale
- ✓ Quando la VS é attuata sotto guida fluoroscopica, l'utilizzo di contrasto deve essere il minore possibile.

Queste chiare linee guida sono state realizzate per aiutare i medici nell'utilizzo dell VS

Appropriateness of clinical and organizational criteria for intra-articular injection therapies in osteoarthritis. A Delphi method consensus initiative among experts in Italy

Marco Paoloni¹, Andrea Bernetti¹, Alberto Beelli², Ovidio Brignoli³, Sandro Buoso⁴, Achille Patrizio Caputi⁵, Fabio Catani⁶, Daniela Coclite⁷, Massimo Fini⁸, Lorenzo Mantovani⁹, Alberto Migliore¹⁰, Antonello Napoletano⁷, Ugo Viora¹¹ and Valter Santilli¹

¹ Dipartimento di Medicina Fisica e Riabilitativa, Sapienza Università di Roma, Rome, Italy

² UOC di Radiologia Diagnostica ed Interventistica, Ospedale San Giovanni Calibita Fatebenefratelli, Rome, Italy

³ Società Italiana di Medicina Generale, Florence, Italy

⁴ UOC di Medicina Fisica e Riabilitativa, ULSS 13 Mirano, Italy

⁵ Dipartimento di Medicina Clinica e Sperimentale Università degli Studi di Messina, Messina, Italy

⁶ Dipartimento di Ortopedia e Traumatologia, Azienda Ospedaliero Universitaria di Modena, Università di Modena e Reggio Emilia, Modena, Italy

⁷ Ufficio Relazioni Esterne, Istituto Superiore di Sanità, Rome, Italy

⁸ Direzione Scientifica, IRCCS San Raffaele Pisana, Rome, Italy

⁹ Centro di Studi e Ricerche sulla Sanità Pubblica, Università degli Studi di Milano Bicocca, Monza, Italy

¹⁰ UOS di Reumatologia, Ospedale San Pietro Fatebenefratelli, Rome, Italy

¹¹ ANMAR – Associazione Nazionale Malati Reumatici Onlus, Italy

“Definizione dei criteri di appropriatezza clinica e organizzativa per il trattamento iniettivo intra-articolare della patologia osteoarticolare degenerativa”

Tabella 1a – Consenso su Accordo	(%)
Terapia iniettiva IA utile nei pazienti con OA del ginocchio e dell'anca	100,0-69,2
Terapia iniettiva IA utile nei pazienti con OA di media gravità	88,5
Il profilo di sicurezza influisce nella scelta del farmaco/medical device	90,4
La rapida azione sui sintomi influisce nella scelta del prodotto	74,5
Il mantenimento a lungo termine dei risultati influisce nella scelta del farmaco/medical device	90,4
L'interazione con altre terapie influisce nella scelta del farmaco/medical device	82,7
Il livello delle evidenze scientifiche influisce nella scelta del farmaco/medical device	86,5
AI ad alto pm e AI reticolato adeguati per terapia iniettiva IA della coxartrosi	82,0-69
AI a medio e alto pm , AI crosslinkato , AI reticolato adeguato per terapia iniettiva IA della gonartrosi	69,2-80

Tabella 1b – Consenso su Accordo	(%)
AI a medio pm con freq. 1-5 monosett. utile per terapia iniettiva IA OA in fase iniziale/di media gravità	68,0
AI reticolato con freq. 1-2 monosett. utile per terapia iniettiva IA OA in fase iniziale/di media gravità	68,0
Mancato utilizzo prodotto adeguato in terapia iniett. IA per OA per problemi di costo/no rimborsabilità	70,0
Eventi avversi gravi sono rari ($\geq 1/10.000$ a $> 1/1.000$)	82,0
Terapia iniettiva IA utile nel management complessivo dell'OA	96,0
Terapia iniettiva IA con AI efficace sui sintomi del paziente con OA	80,0
Terapia iniettiva IA con AI efficace sul controllo oggettivo dell'OA	78,0
Terapia iniettiva IA con AI ad alto pm contribuisce a ritardare/evitare impianto protesi	68,0
Terapia iniettiva IA con AI ad alto pm o con AI crosslinkato contribuisce a ridurre uso FANS e analgesici per via sistemica	80-67

Tabella 1c – Consenso su Accordo	(%)
Studio medico privato setting appropriato per praticare le terapie iniettive IA	81,6
Ambulatorio/poliambulatorio setting appropriato per praticare le terapie iniettive IA	91,8
Struttura ospedaliera/clinica setting appropriato per praticare le terapie iniettive IA	81,6
La guida radiologica/ecografica è utile per effettuare infiltrazioni IA nell'anca	100,0
Terapia iniettiva intra-articolare utile per i pazienti con OA della spalla e della caviglia	69,2
AI a basso pm con freq. 5 monosett. utile per terapia IA di OA iniziale/di media gravità	67,3
AI ad alto pm con freq. 2-3 monosett. utile per terapia IA di OA iniziale/di media gravità	69,2
Eventi avversi lievi dopo terapia IA sono rari ($\geq 1/10.000$ a $< 1/1.000$)	69,2
Terapia iniettiva IA con cortisonici efficace sui sintomi del paziente con OA	78,8

CORTICOSTEROIDI

Intraarticular corticosteroid for treatment of osteoarthritis of the knee (Review)

Bellamy N, Campbell J, Robinson V, Gee T, Bourne R, Wells G

THE COCHRANE
COLLABORATION®

- CS vs placebo: CS superiori a placebo nel **ridurre il dolore e persiste per 2-3 settimane (dopo la differenza non è significativa)**
- **CS vs HA: HA ha una azione più lenta, ma più duratura rispetto a CS**
- "CS vs CS: Triamcinolone esacetone superiore a betametasone

✓ indicazione: fase acuta, versamento

REVIEW

Injectable corticosteroids in treatment of arthritis of the knee

H. Ralph Schumacher, MD,^{a,b} Lan X. Chen, MD, PhD^{a,b}

THE AMERICAN
JOURNAL of
MEDICINE®

Table 1 Potential side effects of intra-articular corticosteroid treatment (and methods that may decrease their incidence)^{8,9,48,57}

Systemic effects (avoid high doses and multiple simultaneous injections, utilize accurate injection techniques)
Tendon weakening/rupture, fat atrophy, muscle wasting, skin pigment changes (avoid misdirected injections)
Septic arthritis (use aseptic technique, withhold therapy in at-risk patients)
Nerve and blood vessel damage (utilize accurate injection techniques)
Postinjection symptom flare/synovitis (avoid same preparation for future injections)
Flushing (avoid high doses)
Anaphylaxis (take careful drug allergy history)
Steroid arthropathy (avoid high doses and overly frequent injections)

- ✓ infiltrazioni CS ripetute possono promuovere la degradazione e la distruzione della cartilagine.
- ✓ Uno studio su modello animale ha evidenziato che i CS IA sono in grado di influenzare negativamente il metabolismo articolare, non solo nella sede della infiltrazione, ma come conseguenza di un effetto sistemico anche a livello delle articolazioni non infiltrate (Celeste et al. 2004).

PRP

The rationale for the use of PRP

Numerous growth factors are within the platelets and exert a specific activity on neoproliferation, on cartilage regeneration and an antiapoptotic effect on chondroblasts:

- - *PDGF regulates the secretion and synthesis of collagen;*
- - *EGF causes cellular proliferation, endothelial chemotaxis and angiogenesis;*
- - *VEGF increases angiogenesis and vascular permeability;*
- - *TGF-beta stimulates the proliferation of undifferentiated MSC, stimulates chemotaxis of endothelial cells and angiogenesis;*
- - *bFGF promotes the growth and differentiation of chondrocytes and osteoblasts, stimulates mitogenesis of mesenchymal cells, chondrocytes and osteoblasts.*

The question is:
have all PRP in each patient at any stage
of disease or age the same quality and
quantity of DGF, EGF, VEGF, TGF-beta
and bFGF ?

We don't know

**At the moment we look only
at the concentration of platelets !**

Factors That the Clinician Should Consider Prior to Using PRP for OA

Younger patients appear to respond better than older patients aged 50 years

Stage of OA (treatment with PRP appears to be more suited for mild OA vs severe)

Optimal concentration of platelets (is still unknown)

Optimal centrifugation method (is still unknown)

Patient platelet count (Low baseline platelet count may preclude injection)

Cellular content (eg, remove leukocytes vs leukocyte rich preparations)

Delivery method (eg, liquid vs spray etc)

Activation method (eg, thrombin vs collagen vs intra-articular activation)

Use of local anesthetic (can have negative effects on chondrocytes)

Image guided vs non-image-guided (US appears to be more optimal)

Frequency of injections and dosing interval (not yet determined)

- ❖ Lack of large and well designed RCTS limits a conclusive statement on the efficacy of PRP
- ❖ it still remains to be determined whether there is only a temporary improvement in symptoms or whether PRP may play a role as disease-modifying therapy.
- ❖ Many clinical questions still have to be clarified: the timing of therapy, the volume and frequency of treatment and the ideal composition of the platelet-rich plasma.

Civinini et al. Clinical Cases in Mineral and Bone Metabolism 2013;

COLLAGENO

868 studies

- Ophthalmology
- Implantology
- Stem cells
- Scaffold
- Dermatology
- Scleroderma
- Neurology
- IA therapy

The screenshot shows a web browser window displaying the ClinicalTrials.gov search results page for the term 'collagen'. The browser's address bar shows the URL: <https://clinicaltrials.gov/ct2/results?term=collagen&pg=1>. The page header includes the ClinicalTrials.gov logo and navigation links such as 'Find Studies', 'About Clinical Studies', 'Submit Studies', 'Resources', and 'About This Site'. A search bar at the top right contains the text 'collagen' and shows '868 studies found for: collagen'. Below the search bar, there are tabs for 'List', 'By Topic', 'On a Map', and 'Search Details'. The 'List' tab is selected, and the page displays a table of search results. The table has columns for 'Rank', 'Status', and 'Study'. The first four results are visible:

Rank	Status	Study
1	Withdrawn	Stem Cells in Wound Healing With Collagen Matrix as a Carrier Condition: Wounds Intervention: Biological: Amniotic Stem Cells and Collagen Matrix
2	Completed	Biological Bone Markers and Hydrolyzed Collagen Supplement in Menopausal Healthy Women Condition: Osteoporosis Interventions: Dietary Supplement: hydrolyzed collagen; Dietary Supplement: wheat protein
3	Completed Has Results	Safety and Efficacy of the Addition of 0.3% Lidocaine With EVOLENCE® Conditions: Aging; Pain Interventions: Device: Evolence; Device: Lidocaine; Drug: topical anesthetic
4	Enrolling by	Simultaneous TransPRK and Corneal Collagen Cross-Linking Condition: Keratoconus

The browser's taskbar at the bottom shows the Start button, several application icons, and the system tray with the date and time '9.16'.

Backgorund

- Recent scientific evidence suggests that injecting treatment with porcine collagen could provide interesting results in improving clinical performances.
- The tropocollagen is the substrate for the reconstruction of collagen fibers and acts as Bio-scaffold

Characteristics of collagen in Guna Collagen Medical Devices			
Type of collagen	Molecular weight of type I tropocollagen	Amount of tropocollagen in 1 vial/2ml	Molecules of tropocollagen in 100 µg of tropocollagen preparation
Type I Tropocollagen	300 kDa	100 µg	2X10¹⁴

TYPE 1 COLLAGEN

$\alpha 2 \beta 1$ -INTEGRIN

FIBROBLAST

PI3K/AKT

NF- κ B

CELL SURVIVAL

*Phosphatidil -inositol-3Kinasis
Mitogen activated protein kinases*

2012 Maria del Nogal

RAS/RAF

ERK1/2

CELL PROLIFERATION

2010 Kuo-Shu Tsai

RAC

MLCK

CYTOSKELETAL ORGANIZATION

1996 ISABELLE MERCIER

RESEARCH ARTICLE

Open Access

A double blind randomized active-controlled clinical trial on the intra-articular use of Md-Knee versus sodium hyaluronate in patients with knee osteoarthritis ("Joint")

Luis Severino Martin Martin¹, Umberto Massafia², Emanuele Bizzi² and Alberto Migliore²

The aim

- to evaluate the use of collagen MD-Knee versus sodium hyaluronate (ARTZ[®]) in patient with knee OA.
- The outcome has been clinically assessed through the OMERACT criteria (Outcome Measures in Rheumatology)

New opportunities

- MD could be used in the failure of VS
- MD may be used also as first line treatment in the management of knee and hip OA
- MD devices may be used for the treatment of patients affected by Chondrocalcinosis.
- The combination of HA and MD collagen could be a potential association to improve the efficacy of IA management of chronic diseases

ISIAT 2017

SAVE THE DATE

Ultrasound-guided intra-articular injections:
practical courses on how to do it

Prague **5-7 october**
2017